

Betulaceae Birch family

Corylus L.

hazel

Jill Barbour and Kenneth A. Brinkman

Ms. Barbour is a germination specialist at USDA Forest Service's National Tree Seed Laboratory, Dry Branch, Georgia; Dr. Brinkman retired from the USDA Forest Service's North Central Forest Experiment Station.

Other common names. Filbert, hazelnut.

Growth habit, occurrence, and use. The hazels *Corylus* L. include about 15 species of large, deciduous shrubs (rarely small trees) that occur in the temperate parts of North America, Europe, and Asia. Some species are grown for their nuts or for ornament, and most species provide food for wildlife. In this country, 4 species have present or potential value for wildlife, shelterbelt, or environmental plantings (table 1). For many years, European hazel has been cultivated for the commercial production of its nuts, known as hazelnuts or filberts, mostly in Europe but to some extent in the United States, especially in the Willamette Valley of Oregon. Years of first cultivation for other species are as follows: American hazel (1798), beaked hazel (1745), and California hazel (1910).

Flowering and fruiting. Male and female flowers are borne separately on 1-year-old lateral twigs of the same plant. They are formed late in the summer and open the following spring before the leaves appear (table 2). The male flowers are borne in clusters of 2 to 5 pendulous catkins, consisting only of stamens. The female flower is budlike, each flower has a single ovary with 2 styles that are strikingly red at pollination (Hora 1981). By late summer or early fall, the fertilized female flowers develop into fruits. These are round or egg-shaped, hard-shelled, brown or dark-tan nuts, each consisting of a single seed, or nut, enclosed in an involucre, or husk, consisting of 2 more or less united hairy bracts (figures 1 and 2). The seeds are naturally dispersed by animals or birds. Large seed crops are produced at irregular intervals, usually every 2 or 3 years (NBV 1946; Vines 1960).

Collection of fruits. Hazelnuts may be eaten by rodents, larger animals, or some birds even before they are fully mature. To reduce such losses, fruits should be picked as soon as the edges of the husks begin to turn brown, which may be as early as mid-August.

Extraction and storage of seeds. The fruits should be spread out in thin layers on wire-mesh screens to dry in a room with high humidity for about 1 month. A Dybig macerator can be used to separate the husk from the hazelnut. The machine is operated without water, and the nuts and husks pour out of the spout (Horvath 1999). An aspirator or screen cleaning machine is then needed to separate the husk debris from the hazelnut. The Westrup brush machine dehisces the hazelnut in a square-wire cylinder and a vacuum sucks out the dust with the seed flowing out the opening in the door (Maloney 1999). Yields and number of seeds per weight vary even within the species (table 3).

Because some dormancy is apparently induced by drying the nuts, hazelnuts were once

thought to be recalcitrant and intolerant of any drying (Hong and Ellis 1996). Recommendations usually were to keep the hazelnuts moist after collection and store them moist over winter (stratification) before planting in the spring (Heit 1967; NBV 1946). Hazelnuts are now considered as orthodox in storage behavior, even though moist storage will prevent deep embryo dormancy for at least several months. Seeds of this genus will also remain viable for a year in unsealed containers at room temperature. Most of the viability of American hazelnut and some of beaked hazelnuts (Brinkman 1974) will be retained if seeds are stored in sealed containers at 5 °C. There are no long-term storage data for hazelnuts.

Pregermination treatments. Newly harvested hazelnuts are not dormant, but inhibitors present in the testa and pericarp are carried to the cotyledons and subsequently through the cotyledonary petioles into the embryonic axis (Bradbeer 1978; Jarvis 1975). Numerous studies have been carried out on the nature of dormancy in European hazel, with most of them concerning the balance of gibberellins and inhibitors and starch synthesis (Arias and others 1976; Bradbeer and Pinnfield 1966; Jarvis 1975; Jarvis and Wilson 1978; Jeavons and Jarvis 1984; Li and Ross 1990). Stratification remains the method used to overcome dormancy, however. Hazel seeds require 2 to 6 months of prechilling before germination will occur (Heit 1968a, b). Three months of cold stratification has proven effective (Dirr and Heuser 1987). Stratification removes the block to gibberellin biosynthesis which begins when the seed is transferred to higher temperatures (Bradbeer and others 1978). In nurseries this can be accomplished by fall sowing or by stratifying outdoors over winter before planting. Seeds may benefit from alternations of warm and cold stratification. Freshly harvested European hazel seeds that were warm stratified for 3 weeks followed by 3 weeks at 4 °C germinated best (Dirr and Heuser, 1987).

Germination tests. Germination is hypogeal (figure 3). The seeds have a dormant embryo and germinate slowly without pretreatment. In one experiment, unstratified seeds of American hazel germinated throughout a year (Brinkman 1974). Gibberillic acid (10^{-4} M) applied to European hazel seeds increased the germination from 64% for the control to 86% at 20 °C (Arias and others 1976). Seeds of this species soaked in ethanol and then 0.1% (w/v) mercuric chloride, when put in a lighted chamber germinated 70% compared to seed germinated in total darkness which was only 9% (Jeavons and Jarvis 1996). Results of limited tests are shown in table 4.

Viability testing by staining the seed with tetrazolium chloride (TZ) is the preferred method of ascertaining the seed's quality (ISTA 1993). Seeds should be cracked and soaked in water for 18 hours. After cutting 1 to 2 mm off the cotyledons at the distal ends and splitting them longitudinally, the embryos should be incubated for 12 to 15 hours in 1% TZ, or 18 to 24 hours in a 0.5% solution. Some unstained tissue is allowed in viable seeds, but interpretation is difficult. Standard germination tests can also be performed once the pericarp is removed and the seeds are prechilled for 2 months at 3 to 5 °C (ISTA 1993).

Nursery practice. Although spring sowing of stratified seeds is feasible, most nurseries plant hazel seeds in the fall (Sus 1925). In Holland, seeds of European hazel are mixed with moist sand for several months before sowing in the fall (NBV 1946). In Tennessee, good results with this species were obtained by storing fresh seed dry at 3 °C until planting in October; average tree percent was 98 based on 80% viability (Zarger 1968). Two seedlots of American hazel planted in November and December gave tree percents of 63 and 48, based on 70 and 60% viability. Seeds of both species were sown 2.5 cm (1 in) deep in drills and covered with 2.5 to 3.75 cm (1 to 1.5 in) of

sawdust. Seedbeds had been fumigated with methyl bromide. If seedling densities are kept low, from 43 to 65/m² (4 to 6/ft²), hazel can be outplanted when 1 year old. European hazel and horticultural varieties are frequently propagated by cuttings, grafting, and tissue culture (Dirr and Heuser 1987).

Hazels are attacked by several fungi. The powdery mildew of hardwoods *Phyllactinia guttata* (Wallr.:Fr.) Lév. (synonym *Phyllactinia corylea* (Pers.) P. Karst.) will defoliate the plant. More serious attacks by the fungal parasite *Nematospora coryli* Peglion cause malformation of the nuts (Hora 1981). Hazelnuts are also attacked by the brown rot of pome and stone fruits *Monilinia fructigena* Honey in Whetzel (synonym *Sclerotinia fructigena* Aderhold. Ex Sacc.) which enters through punctures caused by *Balaninus nuceum*, the nut weevil (Hora 1981).

Literature Cited

- Arias I, Williams PM, Bradbeer JW. 1976. Studies in seed dormancy: 9. The role of gibberellin biosynthesis and the release of bound gibberellin in the post-chilling accumulation of gibberellin in seeds of *Corylus avellana* L. *Planta* 131: 135B139.
- Bradbeer JW, Pinfield NJ. 1966. Studies in seed dormancy: 3. The effects of gibberellin on dormant seeds of *Corylus avellana* L. *New Phytologist* 66: 515B523.
- Bradbeer JW, Arias IE, Nirmala HS. 1978. The role of chilling in the breaking of seed dormancy in *Corylus avellana* L. *Pesticide Science* 9: 184B186.
- Brinkman KA. 1974. *Corylus* L., hazel, filbert. In: Schopmeyer SC, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 343B345.
- Dirr MA, Heuser CW Jr., 1987. The reference manual of woody plant propagation from seed to tissue culture. Athens, GA: Varsity Press. 239 p.
- Fernald ML. 1950. Gray's manual of botany. 8th ed. New York: American Book Co. 1632 p.
- Gorshenin NM. 1941. Agrolesomeliorsiya [in Russian: Agro-forest melioration]. 392 p.
- Heit CE. 1967. Propagation from seed: 11. Storage of deciduous tree and shrub seeds. *American Nurseryman* 126 (10): 12B13, 86B94.
- Heit CE. 1968a. Propagation from seed: 15. Fall planting of shrub seeds for successful seedling production. *American Nurseryman* 128(4): 8B10, 70B80.
- Heit CE. 1968b. Thirty-five years' testing of tree and shrub seed. *Journal of Forestry* 66: 632B634.
- Hong TD, Ellis RH. 1996. A protocol to determine seed storage behavior. Tech. Bull. 1. Rome: International Plant Genetic Resources Institute. 62 p.
- Hora B. 1981. The Oxford encyclopedia of trees of the world. Oxford: Oxford University Press. 288 p.
- Horvath D. 1999. Personal communication. Topeka, IL: Mason State Nursery.
- ISTA [International Seed Testing Association]. 1993. International rules for seed testing. *Seed Science and Technology* 21 (suppl.): 1B288.
- Jarvis BC. 1975. The role of seed parts in the induction of dormancy of hazel (*Corylus avellana* L.). *New Phytologist* 75: 491B494.
- Jarvis BC, Wilson DA. 1978. Factors influencing growth of embryonic axes from dormant seeds of hazel (*Corylus avellana* L.). *Planta* 138: 189B191.

- Jeavons RA, Jarvis BC. 1984. The breaking of dormancy in hazel seed by pretreatment with ethanol and mercuric chloride. *New Phytologist* 96: 551B554.
- Li L, Ross JD. 1990. Starch synthesis during dormancy breakage in oilseed of *Corylus avellana*. *Annals of Botany* 66: 507B512.
- Loiseau J. 1945. *Les arbres et la foret*. 204 p. Paris.
- Maloney J. 1999. Personal communication. Licking, MO: George O. White Nursery.
- Munz PA, Keck DD. 1959. *A California flora*. Berkeley: University of California Press. 1681 p.
- NBV [Nederlandsche Boschbouw Vereeniging]. 1946. *Boomzaden: handleiding inzake het oogsten, behandelen, bewaren en uitzaaien van boomzaden* [in Dutch: Tree seed: handbook on the collection, extraction, storage, and sowing of tree seed]. Wageningen, The Netherlands: Ponsen and Looijen. 171 p.
- Rafn J & Son. [circa] 1928. *Skovfrøkontoret's Frøanalyser gennem 40 Aar, 1887B1927*. Udført paa Statsfrøkontrollen i Köbenhavn. Copenhagen. 5 p.
- Rosendahl CO. 1955. *Trees and shrubs of the Upper Midwest*. Minneapolis: University of Minnesota Press. 411 p.
- Shumilina ZK. 1949. Stratifikatsiya semyan drevesnykh i kustarnikovykh porod. Vsesoyuznoe Nauchno-Issledovatel'skii Institut Agrolesomeliior. Goslestekhzidat, Moskva. [Stratification of seeds of trees and shrubs. Transl. OTS-60-51012, 1961. CFSTI, USDC, NTIS, Springfield, Va. 64 p.].
- Sus NI. 1925. *Pitomnik* [in Russian: The forest nursery]. 27 p.
- Swingle CF, comp. 1939. *Seed propagation of trees, shrubs, and forbs for conservation planting*. SCS-TP-27. Washington, DC: USDA Soil Conservation Service. 198 p.
- Van Dersal WR. 1938. *Native woody plants of the United States: their erosion-control and wildlife values*. Misc. Publ. 303. Washington, DC: U.S. Department of Agriculture. 362 p.
- Vines RA. 1960. *Trees, shrubs, and woody vines of the Southwest*. Austin: University of Texas Press. 1104 p.
- Wappes L. 1932. *Wald und Holz ein Nachschlagebuch für die Praxis der Forstwirte, Holzhändler und Holzindustriellen*. Volume 1. Berlin: J. Neumann. 872 p.
- Zarger TG. 1968. Personal communication (25 September 1968). Tennessee Valley Authority, Division of Forests & Development. [check spell-out, please]

Table 1C *Corylus*, hazel: nomenclature and occurrence

Scientific name & synonym(s)	Common name	Occurrence
<i>C. americana</i> Walt.	American hazel, American filbert	Maine to Saskatchewan, S to Georgia; W to Missouri & Oklahoma
<i>C. avellana</i> L.	European hazel, European filbert, common filbert	Europe, to 1,824 m in central Alps
<i>C. cornuta</i> Marsh. <i>C. rostrata</i> Ait.	beaked hazel, beaked filbert	Newfoundland to British Columbia, S to Georgia, Missouri, & E Colorado
<i>C. cornuta</i> var. <i>california</i> Marsh. (A.DC.) Sharp	California hazel, California filbert	Coast ranges from Santa Cruz N to British Columbia

Source: Brinkman (1974).

Table 2C *Corylus*, hazel: phenology of flowering and fruiting

Species	Location	Flowering dates	Fruit ripening dates
<i>C. americana</i>	C	MarBMay	JulyBSept
<i>C. avellana</i>	Europe	FebBApr	SeptBOct
<i>C. cornuta</i>	Tennessee	JanBFeb	AugBSept
var. <i>californica</i>	California	JanBFeb	Sept-Oct

Sources: Fernald (1950), Loiseau (1945), Munz and Keck (1959), NBV (1946), Rosendahl (1955), Sus (1925), Van Dersal (1938), Vines (1960), Wappes (1932), Zarger (1968).

Table 3 *Corylus*, hazel: seed yield data

Species	Place of collection	Seed wt/fruit wt		Range /kg	No. of cleaned seeds/weight			Samples
		kg/45 kg	lb/100 lb		Average			
					/lb	/kg	/lb	
<i>C. americana</i>	C	11B14	25B30	434B1,623	197B736	1,083	491	11
<i>C. avellana</i>	Europe	27	60	353B1,180	160B535	803	364	244
<i>C. cornuta</i>	C	C	C	937B1,490	425B676	549	249	3
var. <i>californica</i>	California	C	C	882B922	400B418	410	186	C

Sources: Brinkman (1974), Gorshenin (1941), NBV (1946), Rafn (1928), Swingle (1939), Vines (1960), Zarger (1968).

Table 4 *Corylus*, hazel: germination test conditions and results

Species	Medium	Germination test conditions				Germinative energy	Germinative		
		Temp (°C)	Dura- tion	Amt (%)	Period Days		Average (%)	Samples	capacity (%)
<i>C. americana</i>	Sand	30	20	60	10	30	13	2	96
<i>C. avellana</i>	Sand or germinator	30	20	60	C	C	69	13	95
<i>C. cornuta</i>	Sand	30	20	60	1	26	1	1	99
var. <i>californica</i>	Sand	30	20	90	C	C	20	1	62

Sources: Brinkman (1974), NBV (1946), Rafn (1928), Shumilina (1949).